


Euscorpilus

Occasional Publications in Scorpiology



**New Localities of *Iurus dufourei dufourei* (Brullé, 1832)
in the Peloponnese, Greece (Scorpiones: Iuridae)**

Diego Facheris

February 2007 — No. 52

Euscorpilus

Occasional Publications in Scorpiology

EDITOR: Victor Fet, Marshall University, 'fet@marshall.edu'

ASSOCIATE EDITOR: Michael E. Soleglad, 'soleglad@la.znet.com'

Euscorpilus is the first research publication completely devoted to scorpions (Arachnida: Scorpiones). *Euscorpilus* takes advantage of the rapidly evolving medium of quick online publication, at the same time maintaining high research standards for the burgeoning field of scorpion science (scorpiology). *Euscorpilus* is an expedient and viable medium for the publication of serious papers in scorpiology, including (but not limited to): systematics, evolution, ecology, biogeography, and general biology of scorpions. Review papers, descriptions of new taxa, faunistic surveys, lists of museum collections, and book reviews are welcome.

Derivatio Nominis

The name *Euscorpilus* Thorell, 1876 refers to the most common genus of scorpions in the Mediterranean region and southern Europe (family Euscorpilidae).

Euscorpilus is located on Website '<http://www.science.marshall.edu/fet/euscorpilus/>' at Marshall University, Huntington, WV 25755-2510, USA.

The International Code of Zoological Nomenclature (ICZN, 4th Edition, 1999) does not accept online texts as published work (Article 9.8); however, it accepts CD-ROM publications (Article 8). *Euscorpilus* is produced in two *identical* versions: online (ISSN 1536-9307) and CD-ROM (ISSN 1536-9293). Only copies distributed on a CD-ROM from *Euscorpilus* are considered published work in compliance with the ICZN, i.e. for the purposes of new names and new nomenclatural acts. All *Euscorpilus* publications are distributed on a CD-ROM medium to the following museums/libraries:

- **ZR**, Zoological Record, York, UK
- **LC**, Library of Congress, Washington, DC, USA
- **USNM**, United States National Museum of Natural History (Smithsonian Institution), Washington, DC, USA
- **AMNH**, American Museum of Natural History, New York, USA
- **CAS**, California Academy of Sciences, San Francisco, USA
- **FMNH**, Field Museum of Natural History, Chicago, USA
- **MCZ**, Museum of Comparative Zoology, Cambridge, Massachusetts, USA
- **MNHN**, Museum National d'Histoire Naturelle, Paris, France
- **NMW**, Naturhistorisches Museum Wien, Vienna, Austria
- **BMNH**, British Museum of Natural History, London, England, UK
- **MZUC**, Museo Zoologico "La Specola" dell'Universita de Firenze, Florence, Italy
- **ZISP**, Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia
- **WAM**, Western Australian Museum, Perth, Australia
- **NTNU**, Norwegian University of Science and Technology, Trondheim, Norway

New localities of *Iurus dufourei* *dufourei* (Brullé, 1832) in the Peloponnese, Greece (Scorpiones: Iuridae)

Diego Facheris

C/o Museo Civico di Scienze Naturali “E.Caffi”
Piazza Cittadella, 10 - 24129 Bergamo, Italy
email: diego.facheris@alice.it

Summary

New localities of *Iurus dufourei dufourei* (Brullé, 1832) are listed, found during a recent survey in the districts of Achaia, Messinia, and Arcadia in the Peloponnese, Greece. Notes on the habitat and a map are provided.

Introduction

The scorpion genus *Iurus* Thorell is known from southwestern Turkey and southern Greece. In Greece, it is found on the islands of the south Aegean Arch and in the southwestern Peloponnese (Werner, 1902, 1937; Penther, 1906; Vachon, 1948; Kinzelbach, 1975; Francke, 1981; Kritscher, 1993; Crucitti 1995a, 1995b, 1998, 1999a, 1999b; Stathi, 1998; Stathi & Mylonas, 2001; Fet & Braunwalder, 2000; Sissom & Fet, 2000; Parmakelis et al., 2006).

Two subspecies of *Iurus dufourei* (Brullé, 1832) have been described. The nominal subspecies is known from the Peloponnese, Kythira, Gavdos, and Crete. On the other hand, *I. dufourei asiaticus* Birula, 1903 is found in Turkey and the neighboring southern Aegean islands of Karpathos, Kos, Leros, Rodos, Samos, and Megisti (Kastelorizo) (Kinzelbach, 1975; Parmakelis et al., 2006).

The systematics of this genus is still unclear. Most authors who studied it consider *Iurus* a monotypic genus with two subspecies (Kinzelbach, 1975; Kritscher, 1993; Parmakelis et al., 2006) while others distinguished two species (Francke, 1981; Vachon & Kinzelbach, 1987). The DNA data of Parmakelis et al. (2006) seem to confirm the existence of two distinct species (Victor Fet, pers. comm.).

Materials and Methods

The observations were conducted 1–9 August 2005 under stones during the daytime, and with the aid of a UV torch at night. One adult specimen (♀), found near the village of Kato Zachlorou (K. Ζαχλωρού), was

collected and is preserved in 75% alcohol at Museo Civico di Scienze Naturali “E.Caffi”, Bergamo, Italy.

New Localities of *I. dufourei*:

(1) Vorio (Βόρειο), Fig.1, locality 1, 600 m a.s.l., 36°57'30"N, 22°14'19"E on the western slopes of Taygetus: 3 specimens found during the day, under stones in an environment with little surface water;

(2) Artemisia (Αρτεμισία), about 7 km on the road to Kalamata (Καλαμάτα), Fig.1, locality 2, 400 m a.s.l. The locality is in an upper vegetation belt featuring low and sparse vegetation on the roadside embankment and no surface water. On the contrary, near the Nedontas (Νέδωνας) river (a few dozen metres further down) the vegetation is lush in a humid environment. On-the-spot inspections were made both in daytime and at night over the entire area. Only one specimen was found in the upper belt during the night hours, whereas several were seen at the river both in daytime and at night.

(3) Karitena, Kalamiou Monastery (Μ. Καλαμίου), Fig.1, locality 3, 550 m a.s.l., 37°28'52"N, 22°02'25"E. Several specimens found at night on rocks in a humid and shady area.

(4) Kakoureika (Κακουραϊκά), Fig.1, locality 4, 280 m a.s.l., 37°34'46"N, 21°55'14"E. One specimen seen at night on large rock outcrops next to a small stream, 3 km NW of the village near a small church.

(5) Kato Zachlorou (Κ. Ζαχλωρού), Fig.1, locality 5, the Vouraikos (Βουραϊκού) River gorge, 700 m a.s.l., 38°05'27"N, 22°09'11"E. The rocky substrate outcrop in the area of the finding consists of Pleistocene sediments characterised by conglomerate deposits and alluvial boulders in clastic rocks. Overall the rocky mass appears more or less fractured due to the neotectonic




Figure 1: Distribution of *Iurus dufourei dufourei*. White squares, new localities; black circles, localities known from literature.

activity which has lifted the entire area. The sub-vertical valley outline has a typically young morphology produced by the great erosion of the slopes. The specimen was collected during the night near a small spring on the unpaved road 1–2 km from the village.

Description of a Specimen

The single collected female specimen from Kato Zachlorou shows all the characteristics of the nominal subspecies. The body is brown with slightly darker pedipalps and paler legs and vesicle, brown chelicerae which become paler apically without any reticulation. The sternum shape, cheliceral dentition, as well as the distribution and number of trichobothria on the pedipalps correspond with the description given by Francke & Soleglad (1981). Pectinal teeth: 9–9.

Measurements (mm). Total length, 66.0. Carapace length, 9.0; front width, 4.0; rear width, 8.5; distance between rear margin and median eyes, 5.2. Metasomal

segments: segment I length, 4.0; width, 4.1; height 3.4; segment II length, 4.6; width 3.4, height 3.0; segment III length, 4.8; width 3.2, height 3.3; segment IV length, 5.3; width 2.9, height 3.0; segment V length, 9.0; width 2.9, height, 2.7; vesicle (with aculeus) length, 10.0; width 2.8, height 2.4. Pedipalps: femur length, 8.5; width 3.2, height 2.5; patella length, 8.4; width 3.6, height 3.2; chela length, 17.3; width 4.0, height 3.5.

Conclusions

The new data expand the geographic range of *Iurus dufourei dufourei*, which was found for the first time in the northern Peloponnese, Greece. The new locality is more than 60 km away from the previously indicated localities. This survey shows that although *Iurus dufourei* prefers humid and protected places, it can also be found at night with the aid of a UV lamp in open spaces, where solar radiation is strong and direct during the daytime.

Acknowledgments

I would like to thank my mother, Maria Elena, for her precious help during the trip to Greece, as well as Marco Valle, Paolo Pantini, Federico Confortini, and Omar Lodovici of the Museo Civico di Scienze Naturali "E.Caffi" in Bergamo for their essential help in compiling and reviewing this note. I am grateful to Victor Fet and two anonymous reviewers for their comments and suggestions.

References

- BIRULA, A. 1903. Miscellanea Scorpiologica V. Ein Beitrag zur Kenntnis der Skorpionenfauna der Insel Kreta. *Annuaire du Musée zoologique de l'Académie impériale des sciences de St.-Petersbourg*, 8: 295–299.
- CRUCITTI, P. 1995a. *Iurus dufourei* del Peloponneso meridionale: Osservazioni ecologiche e biometriche (Scorpiones, Iuridae). *Bollettino dell'Associazione Romana di entomologia*, 49(3–4) (1994): 1–14.
- CRUCITTI, P. 1995b. *Iurus dufourei* (Brullé) nel Peloponneso occidentale e considerazioni sulla scorfiofauna dei Minthi Óros (Grecia). *Bollettino della Società Entomologica Italiana*, 127 (2): 91–98.
- CRUCITTI, P. 1998. Ricerche bio-ecologiche su *Iurus dufourei* (Brullé, 1832) del Peloponneso sud-occidentale (Scorpiones, Iuridae). *Annali del Museo Civico di Storia Naturale di Ferrara*, 1: 31–43.
- CRUCITTI, P. 1999a. Scorpion species diversity in southwestern Peloponnese, Greece (Scorpiones). *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1: 251–256.
- CRUCITTI, P. 1999b. The scorpions of Anatolia: biogeographical patterns. *Biogeographia*, 20: 81–94.
- FET, V. & M. E. BRAUNWALDER. 2000. The scorpions (Arachnida: Scorpiones) of the Aegean area: current problems in taxonomy and biogeography. *Belgian Journal of Zoology*, 130 (Supplement): 17–22.
- FRANCKE, O. F. 1981. Taxonomic and zoogeographic observations on *Iurus* Thorell (Scorpiones, Iuridae). *Bulletin of the British Arachnological Society*, 5(5): 221–224.
- FRANCKE, O. F. & M. E. SOLEGLAD. 1981. The family Iuridae Thorell (Arachnida, Scorpiones). *Journal of Arachnology*, 9: 233–258.
- KINZELBACH, R. 1975. Die Skorpione der Ägäis. Beiträge zur Systematik, Phylogenie und Biogeographie. *Zoologische Jahrbucher. Abteilung für Systematik, Ökologie und Geographie der Tiere*, 102: 12–50.
- KRITSCHER, E. 1993. Ein Beitrag zur Verbreitung der Skorpione im Östlichen Mittelmeerraum. *Annalen des Naturhistorischen Museums in Wien. Serie B. Botanik und Zoologie*, 94/95, B: 377–391.
- PARMAKELIS, A., I. STATHI, L. SPANOS, C. LOUIS & M. MYLONAS. 2006. Phylogeography of *Iurus dufourei* (Brullé, 1832) (Scorpiones, Iuridae). *Journal of Biogeography*, 33: 251–260.
- PENTHER, A. 1906. Bemerkungen über einige Skorpione aus Kreta. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 56: 60–64.
- SISSOM, W. D. & V. FET. 2000. Family Iuridae. Pp. 409–420 in Fet, V., W. D. Sissom, G. Lowe & M.E. Braunwalder. *Catalog of the Scorpions of the World (1758-1998)*. New York Entomological Society, New York, 690 pp.
- STATHI, I. 1998. *Distribution of scorpions in the central and eastern Mediterranean region and preliminary results on the ecology of the scorpions of Crete*. M. Sc. Thesis, University of Crete, Department of Biology, Irakleio, October 1998.
- STATHI, I. & M. MYLONAS. 2001. New records of scorpion from central and eastern Mediterranean area: biogeographical comments, with special reference to the Greek species. Pp. 287–295 in Fet, V. & P. A. Selden (eds.). *Scorpions 2001. In memoriam Gary A. Polis*. Burnham Beeches, Bucks: British Arachnological Society.
- VACHON, M. 1948. Scorpions récoltés dans l'île de Crète par Mr. le Docteur Otto von Wettstein. *Annalen des Naturhistorischen Museums in Wien*, 56: 60–69.
- VACHON, M. & R. KINZELBACH. 1987. On the taxonomy and distribution of the scorpions of the Middle East. In F. Krupp, W. Schneider & R. Kinzelbach (eds.), *Proceedings of the Symposium on the Fauna and Zoogeography of the Middle East, Mainz, 1985. Beihefte zum Tübinger Atlas des Vorderen Orients, Reihe A (Naturwissenschaften)*, 28: 91–103.

- WERNER, F. 1902. Die Skorpione, Pedipalpen und Solifugen in der zoologisch-vergleichend-anatomischen Sammlung der Universität Wien. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 52: 595–608.
- WERNER, F. 1937. Beiträge zur Kenntnis der Tierwelt der Peloponnes, der Inseln Kytera und Euboea sowie der Kleinen Inseln im Saronischen Golf. I. Reisebericht. IV. Skorpione. *Sitzungsberichte der Akademie der Wissenschaften in Wien. Mathematisch-naturwissenschaftliche Klasse. Abteilung I. Biologie, Mineralogie, Erdkunde*, 146: 135–143.