

9-2015

Lipopolysaccharide induced MAP kinase activation in RAW 264.7 cells attenuated by cerium oxide nanoparticles

Vellaisamy Selvaraj
Marshall University

Niraj Nepa
Marshall University, nepal@marshall.edu


Steven Rogers
Marshall University

Nandini D.P.K. Manne
Marshall University, manne@marshall.edu

Ravi K. Arvapalli
Marshall University, arvapalli@marshall.edu

See next page for additional authors

Follow this and additional works at: http://mds.marshall.edu/sp_psr

 Part of the [Chemicals and Drugs Commons](#), [Medicinal and Pharmaceutical Chemistry Commons](#), and the [Nanomedicine Commons](#)

Recommended Citation

Selvaraj, V., Nepal, N., Rogers, S., Manne, N.D., Arvapalli, R., Rice, K.M., Asano, S., Fankenhanel, E., Ma, J.Y., Shokuhtar, T. and Maheshwari, M. (2015). Lipopolysaccharide induced MAP kinase activation in RAW 264.7 cells attenuated by cerium oxide nanoparticles. *Data in Brief*, 4, 96-99.

This Article is brought to you for free and open access by the Faculty Research at Marshall Digital Scholar. It has been accepted for inclusion in Pharmaceutical Science and Research by an authorized administrator of Marshall Digital Scholar. For more information, please contact zhangj@marshall.edu, martj@marshall.edu.

Authors

Vellaisamy Selvaraj, Niraj Nepa, Steven Rogers, Nandini D.P.K. Manne, Ravi K. Arvapalli, Kevin M. Rice, Shinichi Asano, Erin Fankenhanel, J. Y. Ma, Tolou Shokuhfar, Mani Maheshwari, and Eric R. Blough



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib



Data Article

Lipopolysaccharide induced MAP kinase activation in RAW 264.7 cells attenuated by cerium oxide nanoparticles



Vellaisamy Selvaraj^a, Niraj Nepal^a, Steven Rogers^a,
Nandini D.P.K. Manne^a, Ravikumar Arvapalli^a,
Kevin M. Rice^a, Shinichi Asano^a, Erin Fankenhanel^a,
J.Y. Ma^e, Tolou Shokuhfar^f, Mani Maheshwari^a,
Eric R. Blough^{a,b,c,d,*}

^a Center for Diagnostic Nanosystems, Marshall University, Huntington, WV, USA

^b Department of Pharmacology, Physiology and Toxicology, Joan C. Edwards School of Medicine, Marshall University, Huntington, WV, USA

^c Department of Cardiology, Joan C. Edwards School of Medicine, Marshall University, Huntington, WV, USA

^d Department of Pharmaceutical Sciences and Research, School of Pharmacy, Marshall University, Huntington, WV, USA

^e Health Effects Laboratory Division, NIOSH, Morgantown, WV, USA

^f Department of Mechanical Engineering and Engineering Mechanics, Michigan Technological University, Houghton, MI, USA

ARTICLE INFO

Article history:

Received 24 April 2015

Received in revised form

28 April 2015

Accepted 28 April 2015

Available online 8 May 2015

Keywords:

Sepsis

LPS

MAPK

Raw 264.7 cells

ABSTRACT

High mortality rates are associated with the life threatening disease of sepsis. Improvements in septic patient survivability have failed to materialize with currently available treatments. This article represents data regarding a study published in biomaterials (Vellaisamy et al., Biomaterials, 2015, in press). with the purpose of evaluating whether severe sepsis mortality and associated hepatic dysfunction induced by lipopolysaccharide (LPS) can be prevented by cerium oxide nanoparticles (CeO₂NPs) treatment in male Sprague Dawley rats. Here we provide the information about the method and processing of raw data related to our study publish in Biomaterials and Data in Brief (Vellaisamy et al., Biomaterials, 2015, in press; Vellaisamy et al., Data in Brief, 2015, in press.). The

DOI of original article: <http://dx.doi.org/10.1016/j.biomaterials.2015.04.025>

* Correspondence to: Center for Diagnostic Nanosystems, Room 241N Robert C. Byrd Biotechnology Science Center, 1700 3rd Ave. Marshall University, Huntington, WV 25755-1090, USA. Tel.: +1 304 696 2708; fax: +1 304 696 3766.

E-mail address: blough@marshall.edu (E.R. Blough).

<http://dx.doi.org/10.1016/j.dib.2015.04.022>

2352-3409/© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

data contained in this article evaluates the contribution of MAPK signaling in LPS induced sepsis. Macrophage cells (RAW 264.7) were treated with a range of cerium oxide nanoparticle concentration in the presence and absence of LPS. Immunoblotting was performed on the cell lysates to evaluate the effect of cerium oxide nanoparticle treatment on LPS induced changes in Mitogen Activated Protein Kinases (MAPK) p-38, ERK 1/2, and SAPK/JNK phosphorylation.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table [please fill in right-hand column of the table below]

Subject area	Biology
More specific subject area	Nanomedicine
Type of data	figure
How data was acquired	Cell culture and immunoblotting
Data format	Raw and analyzed
Experimental factors	Raw 264.7 cells culture in the presence of LPS and CeO ₂ nanoparticles
Experimental features	Balanced design that encompasses in vitro observations
Data source location	Huntington, WV USA
Data accessibility	<i>The data presented in this article and is related to [1,2]</i>

Value of the data

- The data can be referenced by other scientists investigating the effects of LPS on RAW 264.7 cells MAPK signaling.
- The data can provide comprehensive analysis of the effect of cerium oxide nanoparticles on LPS-induced MAPK signaling in cell culture
- These data provides a more thorough understanding of the MAPK involvement in LPS-induced macrophage mediated signaling

1. Data, experimental design, materials and methods

1.1. Experimental design and methods

This articles contains data related to the research articles entitled “Inhibition of MAP kinase/NF- κ B mediated signaling and attenuation of lipopolysaccharide induced severe sepsis by cerium oxide nanoparticles in Biomaterials and Data in Brief article entitled “Cerium oxide nanoparticles inhibit lipopolysaccharide induced MAP kinase/NF- κ B mediated severe sepsis” [1,2]. The purpose of this study was to evaluate the contribution of Mitogen Activated Protein Kinase (MAPK) signaling in LPS induced sepsis. For this study, macrophage cells (RAW 264.7) were grown and treated with different concentration of cerium oxide nanoparticle in the presence and absence of LPS. Compared to that seen in the control cells, LPS increased the phosphorylation of p-38, Phosphorylation of ERK 42/44 and Phosphorylation of SAPK/JNK. CeO₂ nanoparticle treatment significantly decreased LPS-induced increases phosphorylation of p-38, Phosphorylation of ERK 42/44 and Phosphorylation of SAPK/JNK.

1.1.1. Macrophage culture and uptake of CeO₂ NPs and protective effective of CeO₂ NPs against LPS

To elucidate the contribution of MAPK signaling in LPS induced macrophage cells, RAW 264.7 cells (ATCC) (1.2×10^5 /ml) were grown in 250 mm culture plate at 30 °C with 5% CO₂ in DMEM high glucose medium supplemented with 5% of fetal bovine, containing 1% Pen/Strep (10,000 U Penicillin




Fig. 1. Effect of CeO₂ nanoparticles on MAPK activation induced by LPS in RAW cells. Cells were exposed to LPS in the presence and absence of CeO₂ nanoparticles for 24 h. Expression of Phospho-p-38 and total p-38, Phospho ERK 42/44 and Phospho SPAK JNK by western blot analysis.

and 10 mg Streptomycin/ml) and serum until 70–80% confluence and the media was replaced with the fresh medium containing different doses of CeO₂ NPs (0, 1, 5, 10, 25, 50, 100 and 1000 ng/ml) for 24 h in the absence and presence of LPS.

1.1.2. Immunoblot analysis of LPS and LPS+CeO₂ nanoparticles treated cells

Cells were washed with cold PBS, collected by scraping and centrifuged at 400 × g for 10 min. Total cell lysates was prepared by cell lysis reagent (Sigma) as outlined by the manufacturer. Protein content was estimated in triplicate using the Bradford reagent with bovine serum albumin as a standard. Western blot was performed as mentioned elsewhere. Fifty μg of total protein per well was then subjected to electrophoresis and transfer to nitrocellulose Hybond-C membranes (Amersham™ Hybond™) using standard conditions. Membranes were incubated overnight at 4 °C with the appropriate primary antibody for Phospho p-38, total p-38, Phospho ERK42/44, Phospho SPAK JNK, Total SPAK JNK (Cell Signaling, Dnavers, MA), washed extensively and then incubated for 1 h at room temperature with a HRP labeled anti-rabbit before detection by ECL (Western Blotting Detection Reagent, GE Health Care Amersham, Piscataway, NJ). Immunoreactive signals were quantified by densitometry using Alpha Innotech software (Santa Clara, California). Beta actin immunoreactivity was used for normalization between samples.

2. Results

Compared to that seen in the control cells, LPS increased the phosphorylation of p-38, Phosphorylation of ERK 42/44 and Phosphorylation of SAPK/JNK. CeO₂ nanoparticle treatment significantly decreased LPS-induced increases phosphorylation of p-38, Phosphorylation of ERK 42/44 and Phosphorylation of SAPK/JNK (Fig. 1).

Conflict of interests

The authors declare that they have no competing interests.

Acknowledgments

This work was supported in part from DOE grant (DE-PS02-09ER-01 to E.R.B).

References

- [1] S. Vellaisamy, N. Nepal, S. Rogers, N. Manne, R. Arvapalli, K. Rice, S. Asano, E. Frankhanel, J. Ma, T. Shokuhfar, et al., Inhibition of MAP kinase/ NF- κ B mediated signaling and attenuation of lipopolysaccharide induced severe sepsis by cerium oxide nanoparticles, *Biomaterials* 59 (2015) 160–171, <http://dx.doi.org/10.1016/j.biomaterials.2015.04.025>. (Epub ahead of print).
- [2] S. Vellaisamy, N. Nepal, S. Rogers, N. Manne, R. Arvapalli, K. Rice, S. Asano, E. Frankenhanel, J. Ma, Shokuhfar, et al., Cerium oxide nanoparticles inhibit lipopolysaccharide induced MAP kinase/NF- κ B mediated severe sepsis, *Data in Brief* (2015). in press.